

# Diaphragmatic Dysfunctions and Their Treatment: Neural Therapy and Manual Medicine as Effective Approaches

Fatma Gülçin URAL NAZLIKUL<sup>1,2,3</sup> and Hüseyin NAZLIKUL<sup>3,4,5,6</sup>

<sup>1</sup>Department of Physical Medicine and Rehabilitation İstanbul Health and Technology University, İstanbul, Turkey

<sup>2</sup>President of the Association for Scientific Complementary Medicine and Regulation (BTR), İstanbul, Turkey

<sup>3</sup>Naturel Health Center, HakkıYeten Cad. No. 23 Kat 3 , Vital Plaza , 34394 İstanbul, Turkey

<sup>4</sup>President of the Manual Medicine Pain Regulation Association (MTAR), İstanbul, Turkey

<sup>5</sup>President of the International Federation of Medical Associations of Neural Therapy (IFMANT) – Badstrasse 3a , CH-3860 Schattenthalb, Switzerland

<sup>6</sup>President of Turkish Scientific Neural Therapy Society and Regulation Therapy (BNR) HakkıYeten Cad. Apt 23/ 14 TR-34394 İstanbul, Turkey

## ABSTRACT

Diaphragmatic dysfunctions are a medical issue whose relevance is often underestimated. They manifest in respiratory and systemic symptoms such as dyspnea, sleep disturbances, chronic fatigue, gastroesophageal reflux, and even heart failure. As the primary respiratory muscle, the diaphragm is pivotal in maintaining trunk stability and regulating intra-abdominal pressure, making its dysfunction capable of exerting profound impacts on overall health.

Causes and Pathophysiology of Diaphragmatic Dysfunctions: Diaphragmatic dysfunctions can arise from a variety of factors, including:

- Chronic stress, which increases tension in the diaphragm and surrounding fascia,
- Myofascial trigger points, which impair muscle contraction and mobility,
- Scar tissue, which limits fascial mobility,
- Trauma, surgeries, or mechanical stress, which may irritate or impair the function of the phrenic nerve.

The phrenic nerve, which innervates the diaphragm, plays a crucial role in the functionality of this muscle. Irritation or damage to the phrenic nerve can significantly impair diaphragmatic movement, leading not only to breathing difficulties but also to dysregulation of the autonomic nervous system and organ functions.

Treatment Approaches: The combination of neural therapy and manual medicine has proven to be particularly effective in treating diaphragmatic dysfunctions. Injections targeting myofascial trigger points and segmental therapy can support diaphragmatic function. Manual diagnostics play a critical role in identifying segmental dysfunctions and fascial adhesions.

**Keywords:** Diaphragmatic dysfunction; Neural therapy; Manual medicine; Phrenic nerve; Myofascial release

## Anatomical Composition of the Diaphragm and the Role of the Phrenic Nerve

The diaphragm is a dome-shaped musculo-tendinous structure that separates the thoracic cavity from the abdominal cavity. Its origins are located in the spine, ribs, and sternum. The central part of the diaphragm comprises a tendon plate (centrum tendineum), while the lateral portions are muscular. The diaphragm's primary function is to contract

during inhalation, expanding the thoracic cavity and allowing air to flow into the lungs. Simultaneously, it exerts pressure on abdominal organs, thereby supporting digestive functions.<sup>[1,2,3,4]</sup>

The diaphragm is primarily innervated by the phrenic nerve, which arises from the cervical spinal roots (C3-C5). Any dysfunction of this nerve—whether due to injury, mechanical blockage, or neurological disorders—can result in diaphragmatic impairment.<sup>[1,5]</sup>


**Citation:** Ural Nazlikul, F.G., Nazlikul, H. Diaphragmatic Dysfunctions and Their Treatment: Neural Therapy and Manual Medicine as Effective Approaches. *Int Clin Med Case Rep Jour.* 2025;4(1):1-8.

**Received Date:** 12 January, 2025; **Accepted Date:** 15 January, 2025; **Published Date:** 17 January, 2025

**DOI:**

**\*Corresponding author:** Hüseyin Nazlikul, Naturel Health Center, Hakkı Yeten Cad. No. 23 Kat 3 , Vital Plaza , 34394 İstanbul, Turkey, President of the Manual Medicine Pain Regulation Association (MTAR), İstanbul, Turkey, President of the International Federation of Medical Associations of Neural Therapy (IFMANT) – Badstrasse 3a , CH- 3860 Schattenthalb, Switzerland, President of Turkish Scientific, **Email:** [hnazlikul@web.de](mailto:hnazlikul@web.de)

**Copyright:** © 2025 by Prof. Dr. Fatma Gülçin Ural Nazlikul and Prof. Dr. Dr. med. Hüseyin Nazlikul. Open Access under the Creative Commons Attribution License.



**Figure 1 :Anatomical illustration of the diaphragm (Diaphragma),**

**Figure 1** depicts an anatomical illustration of the diaphragm (Diaphragma), a central muscle essential for respiration. Below is a detailed description:

**Position and Function:**

- The diaphragm separates the thoracic cavity (thorax) from the abdominal cavity (abdomen). It is a dome-shaped muscle that contracts during inhalation, allowing the lungs to expand <sup>(5,6,7)</sup>.

**Main Components:**

1. Costal arches (Arcuscostalis), which support the diaphragm laterally and anteriorly.
2. The central tendon (Centrum tendineum), the central, non-muscular portion of the diaphragm.
3. Muscle fibers radiating outward from the central tendon and attaching to the ribs, spine, and sternum.
4. The lumbar attachments (Cruradiaphragmatica), which anchor the diaphragm to the spine.
5. The openings that allow essential structures to

1. Sternum
2. Tendon plate of the diaphragm
3. Pars sternalis of the diaphragm
4. xiphoid process
5. Costal arch
6. Pars lumbalis of the diaphragm
7. Pars costalis of the diaphragm
8. Foramen venae cavae


pass through, such as the esophagus (Hiatus oesophageus), the aorta (Hiatus aorticus), and the inferior vena cava (Foramen venae cavae).

6. The nerve supply, primarily via the phrenic nerve (not numbered in the image but anatomically significant), which provides motor and sensory innervation to the diaphragm <sup>(1,5,7)</sup>.

**Functional Importance:**

- The diaphragm is not only crucial for respiration but also contributes to trunk stabilization, regulation of intra-abdominal pressure, and facilitation of blood and lymphatic flow.

Diaphragmatic dysfunctions should not be considered in isolation, as they can have significant clinical consequences, particularly in chronic diseases. Regardless of the underlying cause, diaphragmatic insufficiency can negatively affect sleep, physical performance, and overall well-being.<sup>[5]</sup> Therefore, thorough investigation is essential to determine the exact cause of dysfunction.



1. Xiphoid process of the sternum
2. Tendon plate of the diaphragm (centrum tendineum)
3. Costal arch
4. Passage of the oesophagus (oesophageal hiatus)
5. Aortic slit (hiatus aorticus)
6. Pars lumbalis of the diaphragm
7. 12th rib
8. M. psoas major
9. Lumbar spine
10. Quadratus lumborum muscle
11. Muscle-free area (trigonumlumbosacrale)
12. Pars costalis of the diaphragm
13. Diaphragm muscle (diaphragm)
14. Passage hole for the inferior vena cava
15. Pars sternalis of the diaphragm

**Figure 2: Diaphragm in caudal view**

**Figure 2** illustrates an anatomical depiction of the diaphragm (Diaphragma), a central muscle critical not only for respiration but also for numerous physiological processes. The labeled sections highlight its key anatomical structures and functions.<sup>[5,7]</sup>

### Key Openings in the Diaphragm

The diaphragm contains several essential openings that facilitate the passage of vital structures between the thoracic and abdominal cavities:

- **Hiatus oesophageus:** This opening allows the esophagus to pass through, enabling the transport of food from the throat to the stomach.
- **Hiatus aorticus:** The aorta passes through this opening, distributing oxygen-rich blood from the heart to the rest of the body.
- **Foramen venae cavae:** This passageway accommodates the inferior vena cava, which returns deoxygenated blood from the lower body to the heart.

### Innervation

- The diaphragm is innervated primarily by the phrenic nerve, which provides motor and sensory input. Any impairment or injury to the phrenic nerve can significantly disrupt the diaphragm's functionality, leading to compromised respiratory efficiency.<sup>[1,5,7]</sup>

### Functional Importance

The diaphragm serves multiple essential roles:

- **Respiration:** As the primary respiratory muscle, the diaphragm contracts and flattens during inhalation, increasing the volume of the thoracic cavity and drawing air into the lungs. During exhalation, it relaxes and returns to its dome-shaped resting position, aiding in the expulsion of air.<sup>[5,6,7]</sup>
- **Intra-Abdominal Pressure Regulation:** The diaphragm plays a pivotal role in maintaining intra-abdominal pressure, which is crucial for processes such as digestion, defecation, childbirth, and coughing..<sup>[5,6,7]</sup>
- **Postural Stability:** By working in coordination with the abdominal muscles, the diaphragm contributes to trunk stabilization and overall body posture.
- **Circulatory Support:** The rhythmic movement of the diaphragm facilitates the circulation of blood and lymph between the thoracic and abdominal cavities, supporting cardiovascular and immune system function.<sup>[5,6,7]</sup>

### Clinical Relevance

- Dysfunction of the diaphragm, often resulting from trauma, scar tissue, or chronic stress, can lead to significant clinical symptoms, including respiratory insufficiency, digestive disturbances, and systemic health issues. Effective treatment often involves a combination of Neural Therapy and Manual Medicine, targeting the diaphragm and its associated structures to restore functionality and alleviate symptoms.
- This figure 2 offers a detailed and insightful representation of the diaphragm's anatomy and its essential contributions to human physiology. It underscores the diaphragm's integral role in maintaining overall health and highlights its significance in clinical evaluation and treatment.<sup>[5,6,7]</sup>

### Diaphragmatic Dysfunctions: An Often Overlooked and Underdiagnosed Cause of Health Issues – Effective Treatment through Neural Therapy and Manual Medicine

The diaphragm (diaphragma) is not only the most important respiratory muscle in the body but also plays a central role in core stabilization, digestion, and blood and lymph circulation.<sup>[5,6,7]</sup>

Diaphragmatic dysfunctions are widespread but frequently underdiagnosed or overlooked (ibid.). This is because the symptoms are not always directly attributable to the diaphragm and often manifest as dyspnea, sleep disorders, heart failure, or chronic fatigue.<sup>[6,7,5]</sup>

The diaphragm influences a variety of bodily functions, including respiration, posture, organ and pelvic function, as well as the function of the floor of the mouth. Furthermore, it has a central impact on the cervical spine, the trigeminal system, and the vascular and lymphatic systems. Considering the role of the phrenic nerve is thus essential for the diagnosis and therapy of diaphragmatic dysfunctions.<sup>[1,2,5]</sup>

### Neural Therapy

The injection of local anesthetics into myofascial trigger points, segmental regions, or along the course of the phrenic nerve is considered an effective approach to support diaphragm function. This method can reduce inflammation, normalize muscle tension, and improve blood circulation. It has proven particularly effective in addressing scars and fascial adhesions.<sup>[8,9,10,11,12]</sup>

### Manual Medicine

Manual diagnostics play a crucial role in identifying

dysfunctions. Techniques such as palpation allow for the precise localization of segmental dysfunctions, fascial adhesions, or muscular tension fields.<sup>[12,13,14]</sup> Through manual techniques, fascial mobility is improved, diaphragmatic movement is enhanced, and reflexive control of the nervous system is optimized.<sup>[15]</sup>

The combination of mobilization, myofascial release, and specific breathing exercises supports the long-term restoration of diaphragmatic function.<sup>[15]</sup>

### **Neural Therapy: A Crucial Therapeutic Approach**

Neural therapy is an effective method for treating diaphragmatic dysfunctions. This therapeutic approach involves the injection of local anesthetics such as procaine or lidocaine into specific areas to achieve a regulatory effect on the autonomic nervous system. Procaine and lidocaine not only exhibit analgesic effects but also possess anti-inflammatory, circulation-enhancing, and alkalizing properties. Additionally, they promote nerve fiber regeneration, which can help restore the normal function of the phrenic nerve.<sup>[8,17]</sup> Current studies demonstrate that neural therapy improves diaphragmatic function and provides long-term symptom relief.<sup>[8,16,17]</sup>

### **The Importance of Manual Diagnostics and Mobilization**

A precise and early diagnosis is crucial for the successful treatment of diaphragmatic dysfunctions. In this context, manual diagnostics play a vital role. Through palpation and other manual examination techniques, tensions, asymmetries, and blockages in the diaphragm and adjacent structures can be identified. This diagnostic method offers an effective way to detect functional disorders early and address them in a targeted manner.<sup>[8,18,9]</sup>

In combination with neural therapy, manual mobilization serves as a powerful therapeutic intervention. After neural therapy, techniques such as thoracic spine mobilization, lymphatic drainage, rib articulation, and myofascial release can be employed to enhance diaphragmatic function and support the healing process.<sup>[12,13]</sup>

### **Neural Therapy and Manual Medicine: A Combination for Long-Term Relief**

The integration of neural therapy and manual medicine offers a holistic approach to the treatment of diaphragmatic dysfunctions. While neural therapy regulates the phrenic nerve and the autonomic nervous system, manual medicine addresses mechanical blockages and muscular tension. The combination of

these two approaches results in significant symptom reduction and improved quality of life.<sup>[13,14,15]</sup>

### **Myofascial Release Techniques**

Myofascial release techniques focus on resolving adhesions and tension in connective tissues (fascia), potentially improving mobility. The application of neuromuscular techniques utilizes muscular energy, joint positioning, myofascial or post-isometric relaxation to target soft tissues, including visceral connective and lymphatic tissues.<sup>[19,20]</sup>

### **In Summary:**

Diaphragmatic dysfunctions are often insufficiently considered during diagnostic evaluations, despite their potential to cause severe symptoms such as dyspnea, heart failure, and chronic fatigue.<sup>[21,19]</sup> The phrenic nerve, as the primary nerve of the diaphragm, plays a central role, and its dysfunction can significantly impair diaphragmatic mobility.

Neural therapy, particularly using procaine and lidocaine, has proven to be an effective method for treating diaphragmatic dysfunctions. It exhibits anti-inflammatory, analgesic, and circulation-enhancing effects.<sup>[22,23]</sup>

Manual diagnostics and mobilization are crucial for the early detection and treatment of diaphragmatic dysfunctions. When combined with neural therapy, they offer an effective therapeutic approach. A holistic approach can lead to significant symptom reduction and sustainable relief.

### **CONCLUSIONS**

Breathing is a systemic activity involving various parts of the body. The health of the diaphragm is critical not only for patients with respiratory conditions but also for those with a wide range of other clinical issues. Restrictions in diaphragmatic mobility, whether during inhalation, exhalation, or within the thoracic region, can result in significant discomfort and dysfunction.

Effective training and targeted treatment of the diaphragm can be advantageous in many clinical scenarios. However, the manual diagnosis of diaphragmatic dysfunctions has been described by only a few authors. This article hypothesizes a diagnostic approach for diaphragmatic disorders using invasive methods and describes a combined therapy involving neural therapy and manual medicine as an effective treatment strategy.

The combination of neural therapy and manual medicine thus represents a highly effective treatment option. Manual medical examinations, supported

by the use of ultrasound, provide a simple and effective method for early detection and treatment of diaphragmatic dysfunctions. In chronic diseases, the possibility of a diaphragmatic dysfunction should always be considered.

Early diagnosis and treatment using this combination therapy can lead to significant improvements in patients' quality of life and prevent disease progression.

**Declaration of conflicts of interest:** The authors

have no conflicts of interest to declare.s

**Sources of funding:** This research received no specific funding from public, commercial or non-profit organisations.

No human or animal studies were conducted by the authors for this paper.

**Author contributions:** The contributions of the authors Nazlikul, H. and Ural Nazlikul should be regarded as equivalent.

## REFERENCES

1. Clara M. Das Nervensystem des Menschen, Verlag Von Johann Ambrosius Barth, Leipzig 1942.
2. Cohen AJ, Katz MG, Katz R, Mayerfeld D, Hauptman E, Schachner A. Phrenic nerve injury after coronary artery grafting: Is it always benign? *Ann. Thorac. Surg. Ann Thorac Surg.* 1997;64(1):148-53.
3. AnatRatnovsky, David Elad. Anatomical model of the human trunk for analysis of respiratory muscles mechanics. *RespirPhysiolNeurobiol.* 2005;148(3):245-62.
4. Nazlikul H, Elmacioglu MA, Ural Nazlikul FG. As Part of a Fictitious Consideration of the Potential Effects of the 2019 Coronavirus Pandemic (SARS), it is Hypothesised that Neural Therapy with local Anaesthetics Could be Able to Mitigate the Adverse Effects Associated with the Pandemic and its Post-SARS Phase. *J Pulm Med Respir Res.* 2024;10:092.
5. Nazlikul H. Diaphragma-Dysfunktionen: Eine oft übersehene Ursache von Gesundheitsproblemen – Effektive Behandlung durch Neuraltherapie und Manuelle Medizin. *Ganzheitsmedizin Heft 4, Jahrgang 37, Dezember 2024.*
6. Anraku M, Shargall Y. Surgical conditions of the diaphragm: Anatomy and physiology. *Thorac. Surg. Clin. ThoracSurgClin.* 2009;19(4):419-29.
7. Anatomie J. Das Zwerchfell: zwei physiologische Muskeln in einem. *J Anat.* 2002;201(4):305-312.
8. Nazlikul H. Nöralterapi Ders Kitabı, 2. Baskı, İstanbul Nobel Tıp Kitapevi 2015 - Neuraltherapy Textbook, 2nd Edition, İstanbul Nobel Medical Bookstore 2015.
9. Fischer L. Physical and neurobiological principles. In: Liem, T., van den Heede, P. (Eds.), *Foundations of Morphodynamics in Osteopathy.* Handspring Publishing, Edinburgh. 2015;67-96.
10. Fischer L, Barop H, Maxison-Bergemann S. Health Technology Assessment HTA Neural therapy according to Huneke. Program Evaluation Complementary Medicine (PEK). On behalf of the Swiss Federal Office of Public Health.
11. Fischer L, Ludin SM, Thommen D, Hausammann R. Application for the Adoption of Interference Field Therapy (Neural Therapy According to Huneke) Into the Health Benefit Basket of the Compulsory Health Insurance. Submitted to the Swiss Federal Office of Public Health. 2010.
12. Nazlikul H, Ural Nazlikul FG. Regülasyontıbbındanöralterapi vemanuelterapiuygulamalarınınyeri = The Place of Neural Therapy and Manual Therapy Applications in Regulatory Medicine. Türk A, editör. *Regülasyontıbbındasporvesağlık.* 1. baskı. Ankara: Türkiye Klinikleri; 2023;8-14.
13. Nazlikul H. Thorakale Blockaden und Wirkung der Neuraltherapie in Kombination mit manueller Therapie, *Manuelle Medizin.* 2010;48:329-338.
14. Nazlikul H, Ural Nazlikul FG. Diagnostik und Therapie der ersten Rippe: eine neue Technik. *Manuelle Medizin.* 2015;1-6.
15. Nazlikul H. Die segmentale vertebrale Dysfunktion ist ein multikausales Geschehen. *Manuelle Medizin.* 2014;52(5):432-6.
16. Fischer L. Neuraltherapie: Neurophysiologie, Injektionstechnik und Therapievorschläge, 5. Aufl. Thieme, Stuttgart.
17. Fischer L. Stellate ganglion. In: Weinschenk, S. (Ed.), *Handbuch Neuraltherapie – Diagnostik und Therapie mit Lokalanästhetika.* Thieme, Stuttgart. 2020;413-416.
18. Türk A, Melekoğlu T, Nazlikul H. Chronic Pain and Physical Activity = Kronik ağrı ve fiziksel aktivite. Kavukcu E, editör. *KronikHastalıkların-*

Tedavisinde Spor ve Fiziksel Aktivitenin Yeri =  
The Place of Sports and Physical Activity in the  
Treatment of Chronic Diseases. 1. Baskı. Ankara:  
Türkiye Klinikleri; 2022. p.1-8.

19. Ural Nazlıkul FG, Bölükşenlikci H. Lökomotor-  
sistem hastalıklarından nöralterapi. Babacan CA,  
editör. Ozon ve Nöralterapi. 1. Baskı. Ankara:  
Türkiye Klinikleri. 2022;61-6.

20. Vinyes D, Muñoz-Sellar M, Albareda Colilles  
G, Gurevich ML. Procaine injections in myo-  
fascial tension points in the treatment of anxiety  
disorders: A case series. International Journal of  
Clinical Case Reports and Reviews. 2025;22(1).

21. Böhni UW, Lauper M, Locher H. Manuelle  
Medizin 2. Diagnostische und therapeutische  
Techniken praktisch anwenden. Stuttgart,  
Thieme. 2012.

22. Gold-Szklarski K, Ortner W, Osztovcics J. Neu-  
raltherapie-Grundlagen und Anwendung. In:  
Frass, M., Krenner, L. (eds) Integrative Medizin.  
Springer, Berlin, Heidelberg. 2019.

23. Fischer L, Barop H, Ludin SM, Schaible HG.  
Regulation of acute reflectory hyperinflammation  
in viral and other diseases by means of stellate  
ganglion block. A conceptual view with a focus  
on Covid-19. Auton Neurosci. 2022;237:102903.